

**Table 1**  
**Formulations Used in Excipient Compatibility Study**

No	Test Excipient (%)	Other Ingredients (%)
F1	Control	Avicel PH 112 (77), Explotab (8), Syloid (0.25), Mg Stearate (0.25)
F2	Talc (0.25)	Avicel PH 112 (77), Explotab (8), Mg Stearate (0.25)
F3	Stearic Acid (1)	Avicel PH 112 (76), Explotab (8), Syloid (0.25)
F4	Crospovidone (5)	Avicel PH 112 (80), Syloid (0.25), Mg Stearate (0.25)
F5	Nu-Tab (77)	Explotab (8), Syloid (0.25), Mg. Stearate (0.25)
F6	Avicel PH 112 (38.6), Nu-Tab (38.6)	Explotab (8), Syloid (0.25), Mg. Stearate (0.25)
F7	Nu-Tab (39.9), Crospovidone (5)	Avicel PH 112 (39.9), Syloid (0.25), Mg. Stearate (0.25)
F8	Nu-Tab (40), Crospovidone (5), no Syloid	Avicel PH 112 (40), Mg. Stearate (0.5)

**Table 2**  
**Formulations Investigated to Select Anti-Oxidant**

No.	Antioxidant (%)	Other Excipient	Tablet Wt (mg)
W1	None	Crospovidone	200
W2	Methionine (0.5)	Crospovidone	20
W3	Ascorbic Acid (1)	Crospovidone	200
W4	Methionine (0.5)	None	250
W5	Methionine (0.5) (EDTA) (0.8)	None	250

**Table 3**  
**rhIL-11 Leading Tablet Formulation**  
 Manufactured by Fluid Bed Granulation

<b>Ingredients</b>	<b>mg / tablet</b>
<i>Intragranular</i>	
rhIL-11 (concentrate equivalent to 2.5 mg)	5.561
Avicel PH 102	92.50
Na <sub>2</sub> HPO <sub>4</sub> Anhydrous	8.50
NaH <sub>2</sub> PO <sub>4</sub> Anhydrous	6.50
Methionine	1.00
Tween 80	0.339
<i>Extragranular</i>	
Avicel PH 112	73.5
Na <sub>2</sub> HPO <sub>4</sub> Anhydrous	4.25
NaH <sub>2</sub> PO <sub>4</sub> Anhydrous	3.25
Explotab	4.00
Magnesium Stearate	0.60
Total	200
Coating	
Eudragit L30D	5%

**Table 4**  
**Effect of Physical Stress on the Integrity of**  
**rhIL-11**

Hardness (Kp)	Recovery <sup>a</sup> (%)	Multimer <sup>b</sup> (%)	Met 58 (%)	Related (%)
2.4	111.0	0.2	4.1	3.7
4.0	105.3	0.3	4.2	3.9
7.5	96.4	0.3	4.4	4.1
12.8	100.2	0.2	4.3	4.0

<sup>a</sup> Measured by RP-HPLC <sup>b</sup> measured by Size  
Exclusion Chromatography.

**Table 5**  
***In Vitro* Bio-activity by T-10 bioassay**  
 (Directly compressed tablets of rhIL-11)

<b>Formulation</b>	<b>Sp Act Uwho/mg</b>	<b>IC Sp Act Uwho/mg</b>
Tablet: Crospovidone, Syloid, Avicel, Mg Stearate	5.82E+06	6.70E+06
Blend: Avicel, Nu- Tab, Explotab, Syloid, Mg Stearate	6.57E+06	5.80E+06
Tablet: Avicel, Nu- Tab, Explotab, Syloid, Mg Stearate	6.38E+06	7.70E+06

**Sp Act: Specific Activity; IC Sp Act: Internal Control  
Specific Activity**

**Table 6**  
**Stability of Enteric Coated Tablets of rhIL-11**  
**(by Fluid Bed Granulation)**

Time (Weeks) (Conditions)	Strength (%)	Met <sup>58</sup> (%)	Related Species (%)
Initial	93.6	5.0	6.7
2 (40°C/75%RH)	86.9	4.5	3.4
4 (40°C/75%RH)	86.6	5.0	3.8
15 (Room Temp.)	94.1	4.0	4.9

**Table 7: Sustained Release Tablet  
Formulations Prepared by Direct Compression**

Ingredients	Formulation 1 (%)	Formulation 2 (%)	Formulation 3 (%)
Lyophilized rhIL-11*	6.3	6.0	5.7
HPMC (Methocel K4M PREM)	10.5	15	19
Microcrystalline Cellulose (Avicel PH112)	10.5	10	9.5
Sucrose (NU-TAB®)	68.5	65	62
Silicon Dioxide (Syloid)	0.26	0.25	0.24
Mg-stearate	0.79	0.75	0.71
Na <sub>2</sub> HPO <sub>4</sub> (Anhydrous)	1.78	1.7	1.62
NaH <sub>2</sub> PO <sub>4</sub> (Anhydrous)	1.37	1.3	1.24

\* Each tablet contains 2.5 mg rhIL-11.

**Table 8: Composition of Sustained Release  
Tablet Formulations Prepared by High Shear  
Wet Granulation**

Ingredients	Formulation 4 (%)	Formulation 5 (%)
rhIL-11*	1.0	1.0
Methocel K4M PREM	10.0	15.0
Avicel PH112	30.0	30.0
NU-TAB®	55.04	50.04
Syloid	0.25	0.25
Mg-stearate	0.74	0.74
Na <sub>2</sub> HPO <sub>4</sub> (Anhydrous)	1.68	1.68
NaH <sub>2</sub> PO <sub>4</sub> (Anhydrous)	1.29	1.29

\* Each tablet contains 2.5 mg rhIL-11 added  
as bulk solution.



**Table 9: Composition of Sustained Release  
Tablet Formulations Prepared by Fluid Bed  
Granulation Using Higher Viscosity Grades of  
HPMC**

Ingredients	Formulation	Formulation	Formulation
	6	7	8
	(%)	(%)	(%)
rhIL-11 Granules*	48.6	45.7	45.7
Methocel K4M PREM	31.9	25	24
Methocel K15M PREM	-----	5.3	6.0
Mannitol	18.44	23.0	15.3
Avicel PH102	-----	-----	8.0
Syloid	0.26	0.25	0.25
Mg-Stearate	0.8	0.75	0.75

\* Prepared by fluid bed granulation. Equivalent to 2.5 mg rhIL-11 per tablet.

**Table 10: Composition of Sustained Release  
Tablet Formulations Prepared by Fluid Bed  
Granulation Using Lower Viscosity Grades of  
HPMC and Various Phosphate Buffer Species**

Ingredients	Formulation 9 (%)	Formulation 10 (%)	Formulation 11 (%)	Formulation 12 (%)
rhIL-11 Granules*	45.7	45.7	45.7	45.7
Methocel K100 LV, LH, CR, Premium	25.0	30.0	25	25
Mannitol	16.3	-----	-----	28.3
Syloid	0.25	0.25	0.25	0.25
Mg-Stearate	0.75	0.75	0.75	0.75
Na <sub>2</sub> HPO <sub>4</sub>	6.8	13.3	-----	-----
NaH <sub>2</sub> PO <sub>4</sub>	5.2	10	-----	-----
(NH <sub>4</sub> ) <sub>2</sub> HPO <sub>4</sub>	-----	-----	16.1	-----
(NH <sub>4</sub> )H <sub>2</sub> PO <sub>4</sub>	-----	-----	12.2	-----

\*Prepared by fluid bed granulation. Equivalent to 2.5  
mg rhIL-11

**Table 11: Composition of IL-11 Delayed Release Multiparticulate Capsules**

Component	Percentage (% wt/wt)	Target for 5 mg Capsul (mg)
rhIL-11	1.10 <sup>b</sup>	5.500
Sugar spheres, NF	68.0	339.9
Glycine, USP	2.47	12.38
Sodium phosphate (dibasic), USP	0.180	0.8855
Sodium phosphate (monobasic), USP	0.060	0.3037
Polysorbate-80, NF	0.028	0.1377
Methionine, USP	0.206	1.028
Hydroxypropyl methylcellulose, USP	3.91	19.57
Methacrylic acid copolymer dispersion, NF (Eudragit L30D-55)	15.0	74.95
Talc, USP	7.50	37.49
Sodium hydroxide, NF	0.090	0.4496
Triethyl citrate, NF	1.50	7.490
Purified water, USP	Removed during processing	q.s.
Size #0 Hard gelatin capsule		
Total		500 mg

A 10% overage rhIL-11 is used to compensate for losses during manufacture.

Label/Package
---------------

**Table 12 rhIL-11 Delayed Release Capsules, 5 mg/Capsule  
Long Term Storage at 2-8°C, 0-18 Months**

Tests	Impurities &							
	Strength	Total Inactive Species	Met <sup>58</sup> – Oxidized Species	rhIL-11 Related Species	Specific Activity (T-10 Bioassay)	Dissolution - Acid		Moisture
						Stage (0.1 N HCl)	Dissolution Buffer Stage (Phosphate Buffer)	
Initial	4.60	9.4 %	6.7 %	2.7 %	8.1 x 10 <sup>6</sup>	3 %	76 %	1.1 %
1 Month	4.94	7.1 %	4.5 %	2.7 %	NS <sup>b</sup>	3 %	74 %	1.3 %
83 Days	4.94	6.3 %	4.0 %	2.3 %	7.0 x 10 <sup>6</sup>	3 %	82 %	1.1 %
6 Months	4.74	7.3 %	4.4 %	3.0 %	7.0 x 10 <sup>6</sup>	2 %	84 %	1.1 %
9 Months	5.02	8.1 %	5.3 %	2.8 %	1.1 x 10 <sup>7</sup>	3 %	65 %	2.4 %
12 Months	4.49	5.0 %	3.2 %	1.9 %	8.9 x 10 <sup>6</sup>	NT	NT	1.6 %
18 Months	4.60	5.8 %	4.0 %	1.8 %	8.0 x 10 <sup>6</sup>	1 %	69 %	1.1 %

**Table 13**      **rhIL-11 Delayed Release Capsules, 5 mg/Capsule**  
**Long Term Storage at 25°C, 0-18 Months**

Tests	Strength	Impurities &							
		Total		rhIL-11			Dissolution - Acid Stage (0.1 N HCl)	Dissolution Buffer Stage (Phosphate Buffer)	Moisture
		Inactive Species	Met <sup>58</sup> – Oxidized Species	Related Species	Specific Activity (T-10 Bioassay)				
						Average			
Initial	4.60	9.4 %	6.7 %	2.7 %	8.1 x 10 <sup>6</sup>	3 %	76 %	1.1 %	
1 Month	4.86	7.3 %	4.6 %	2.7 %	NS <sup>b</sup>	2 %	76 %	1.4 %	
83 Days	4.82	6.6 %	4.0 %	2.5 %	6.9 x 10 <sup>6</sup>	3 %	80 %	1.2 %	
6 Months	4.75	9.1 %	5.4 %	3.7 %	5.7 x 10 <sup>6</sup>	1 %	75 %	1.2 %	
9 Months	4.87	10.3 %	6.7 %	3.6 %	1.2 x 10 <sup>7</sup>	2 %	65 %	1.5 %	
12 Months	4.48	7.9 %	5.1 %	2.9 %	7.4 x 10 <sup>6</sup>	2 %	68 %	2.1 %	